

Native Landscape Seed Mix*

Used For All Non-Sodded Areas of The WSSI Landscape

| <u>Common Name</u> | <u>Scientific Name</u> |
|----------------------------|--------------------------------------|
| Grain Rye | <i>Secale cereale</i> |
| Annual Ryegrass | <i>Lolium multiflorum</i> |
| Silky Wild Rye | <i>Elymus villosus</i> |
| Camper Little Bluestem | <i>Andropogon scoparius</i> 'Camper' |
| Indian Grass | <i>Sorghastrum nutans</i> |
| Deer Tongue | <i>Panicum clandestinum</i> |
| Purple Top | <i>Tridens flavus</i> |
| Bottlebrush Grass | <i>Hystrix patula</i> |
| Common Broom Sedge | <i>Andropogon virginicus</i> |
| Black Eyed Susan | <i>Rudbeckia hirta</i> |
| Partridge Pea | <i>Chamaecrista fasciculata</i> |
| Lance Leaved Coreopsis | <i>Coreopsis lanceolata</i> |
| Wild Senna | <i>Senna hebecarpa</i> |
| Marsh Blazing Star | <i>Liatris spicata</i> |
| Tall White Beardstongue | <i>Penstemon digitalis</i> |
| Blue False Indigo | <i>Baptisia australis</i> |
| Virginia Spiderwort | <i>Tradescantia virginiana</i> |
| Canada Anemone | <i>Anemone canadensis</i> |
| Butterfly Milkweed | <i>Asclepias tuberosa</i> |
| Appalachian Beard Tongue | <i>Penstemon laevigatus</i> |
| New England Aster | <i>Aster novae-angliae</i> |
| Wild Quinine | <i>Parthenium integrifolium</i> |
| Whorled Rosinweed | <i>Silphium trifoliatum</i> |
| Joe Pye Weed | <i>Eupatoriadelphus fistulosum</i> |
| Spotted Beebalm | <i>Monarda punctata</i> |
| Eastern Columbine | <i>Aquilegia canadensis</i> |
| Zigzag Aster | <i>Aster prenanthoides</i> |
| Licorice Scented Goldenrod | <i>Solidago odora</i> |
| Grass Leaved Goldenrod | <i>Euthamia graminifolia</i> |

*Plants listed in order of highest percentage to lowest percentage of seed in the mix.